

Congratulations on Your Pregnancy

Included in this pack is important information to help you have a healthy pregnancy as well as information on how to arrange for your antenatal care.

NEW MATERNITY SELF-BOOKING

The Maternity Digital Transformation Programme launched a new online self-referral form for newly pregnant women to register for maternity services. The form collects basic demographic details alongside basic obstetric and medical history. Once a self-referral has been submitted a midwife will collect you directly to offer early pregnancy advice and to arrange an appointment at a suitable time and place. Please complete the online form to allow direct communication with your midwife before your appointment.

<https://www.badgernotes.net/selfreferral/carelocation/somerset>

If you have no internet access, please complete the form titled 'Congratulations on your pregnancy' and return this to reception. This will then be passed to the midwife who will be in contact to book your first appointment with them. This first appointment should ideally be when you are 10 weeks pregnant so that your first scan is around 12-14 weeks.

Second, please book a routine appointment to see your GP.

Third, please read the leaflet which gives you useful information on how to have a healthy pregnancy. This can also be found at: <https://patient.info/health/planning-to-become-pregnant>

Finally, just to let you know we will write to you when you are around 16 weeks pregnant to offer you a whooping cough vaccine, to protect your baby from this when he or she is born. If you are pregnant during the winter months, we also recommend you have a flu vaccine as you are at more risk of complications if you do get the flu whilst pregnant.


Pregnancy Self Referrals

When a lady becomes pregnant, please could they complete the Online Self-Referral to Badgernotes. The old Online Referral Service is no longer available.

The new link for this service is:

Badger Notes – Self-refer your pregnancy to Somerset

www.badgernotes.net/SelfReferral/CareLocation/somerset



Planning your pregnancy

You can improve your chances of getting pregnant and having a healthy pregnancy by following the steps on this page.

Take a folic acid supplement

It's recommended that you should take a daily supplement of folic acid when you're pregnant, or there's a chance you might get pregnant.

You should take a 400 microgram supplement of folic acid every day before you get pregnant, and every day afterwards, up until you're 12 weeks pregnant.

A microgram is 1,000 times smaller than a milligram (mg). The word microgram is sometimes written with the Greek symbol μ followed by the letter g (μg).

Folic acid reduces the risk of your baby having a neural tube defect, such as [spina bifida](#).

A neural tube defect is when the foetus's spinal cord (part of the body's nervous system) does not form normally.

You might be advised to take a higher dose supplement of 5 milligram (5mg) every day.

You may need to take a 5mg supplement of folic acid if:

- you or the baby's other biological parent have a neural tube defect
- you previously had a pregnancy affected by a neural tube defect
- you or the baby's other biological parent have a family history of neural tube defects
- you have diabetes

Your doctor may also recommend a 5mg supplement if you take certain medicines, such as anti-epilepsy medicine or anti-retroviral medicine for HIV.

Talk to a GP if you think you need a 5mg dose of folic acid, as they can prescribe a higher dose.

You can get folic acid tablets at pharmacies, or talk to a GP about getting a prescription.

Do not worry if you get pregnant unexpectedly and were not taking a folic acid supplement at the time. Start taking them as soon as you find out, until you're past the first 12 weeks of pregnancy.

Stop smoking

[Smoking during pregnancy](#) has been linked to a variety of health problems, including:

- premature birth
- low birth weight
- [sudden infant death syndrome \(SIDS\)](#), also known as cot death
- [miscarriage](#)

Quitting can be hard, no matter how much you want to, but support is available.

[NHS Quit Smoking](#) offers free help, support and advice on stopping smoking, including when you're pregnant, and can give you details of local support services.

Smoke from other people's cigarettes can damage your baby, so ask your partner, friends and family not to smoke near you.

Cut out alcohol

Do not drink alcohol if you're pregnant or trying to get pregnant. Alcohol can be passed to your unborn baby.

Drinking in pregnancy can lead to long-term harm to your baby, and the more you drink, the greater the risk.

Find out about [alcohol and pregnancy](#).

Keep to a healthy weight

If you're overweight, you may have problems getting pregnant and fertility treatment is less likely to work.

Being overweight (having a BMI over 25) or obese (having a BMI over 30) also raises the risk of some pregnancy problems, such as [high blood pressure](#), [deep vein thrombosis](#), [miscarriage](#) and [gestational diabetes](#).

Before you get pregnant you can calculate your BMI. But this may not be accurate once you're pregnant, so consult your midwife or doctor.

Having a healthy diet and doing moderate exercise are advised in pregnancy, and it's important not to gain too much weight.

You can keep to a healthy weight by having a [balanced diet](#) and doing regular exercise. [Read more about exercise in pregnancy](#)

Know which medicines you can take.

Not all medicines are safe to take when you're pregnant or planning a pregnancy, whether they're on prescription or medicines you can buy in a pharmacy or shop.

Information:

If you take prescribed medicine and you're planning to get pregnant, talk to a doctor.

Do not stop taking your medicine without talking to a doctor.

[Find out about medicines in pregnancy.](#)

Vaccinations and infections

Some infections, such as [rubella \(German measles\)](#), can harm your baby if you catch them during pregnancy.

Most people in the UK are immune to rubella, thanks to the uptake of the [measles, mumps and rubella \(MMR\) vaccination](#).

If you have not had 2 doses of the MMR vaccine, or you're not sure if you have, ask your GP surgery to check your vaccination history.

If you have not had both doses or there's no record available, you can have the vaccinations at your GP surgery.

You should avoid getting pregnant for 1 month after having the MMR vaccination, which means you'll need a reliable method of [contraception](#).

Talk to a doctor if you have a long-term condition

If you have a long-term condition, such as epilepsy or diabetes, it could affect the decisions you make about your pregnancy – for example, where you might want to give birth.

Before you get pregnant, have a discussion with your specialist or a GP about getting pregnant.

You can also talk to your GP if you have a mental health condition and are planning to get pregnant. They may be able to refer you to a specialist mental health team who offer support before, during and after pregnancy.

If you're taking medicine for a condition, do not stop taking it without talking to a doctor.

Testing for sickle cell and thalassaemia

Sickle cell disease (SCD) and thalassaemia are inherited blood disorders that mainly affect people whose ancestors come from Africa, the Caribbean, the Mediterranean, India, Pakistan, south and Southeast Asia, and the Middle East.

If you are pregnant and live in England you will be offered screening tests for these disorders, but you do not have to wait until you're pregnant before you have a test.

If you or your partner are concerned you may be a carrier for 1 of these disorders, perhaps because someone in your family has a blood disorder or is a carrier, it's a good idea to get tested before starting a family.

You can ask for a free blood test from either a GP or a local sickle cell and thalassaemia centre.

Find out more about [screening for sickle cell and thalassaemia in pregnancy](#)

More in [Trying for a baby](#)

- [Trying to get pregnant](#)
- [Getting pregnant if you have a mental health condition](#)
- Planning your pregnancy
 - [Doing a pregnancy test](#)
 - [Signs and symptoms of pregnancy](#)
 - [How long does it usually take to get pregnant?](#)
 - [Planning another pregnancy](#)

Page last reviewed: 26 April 2023
Next review due: 26 April 2026

Planning to become pregnant

Reviewed by [Dr Colin Tidy, MRCGP](#). Last updated by [Dr Hayley Willacy, FRCGP](#). Last updated 28 Apr 2020
Meets Patient's [editorial guidelines](#)

This leaflet offers advice if you are planning to become pregnant. The aim is to plan ahead to reduce the risk of harm to the baby and to reduce the risk of pregnancy complications. If you take any regular medication, or have a long-term medical condition, see your doctor to discuss any changes which may need to be made. Do this before you are pregnant ideally, in case anything might affect your baby.

There is a summary checklist at the end of the leaflet.

If you are already pregnant, you may find [Diet and Lifestyle during Pregnancy](#) more help.

When should I have my next baby?

A recent study showed that mothers should wait for 12 to 18 months after giving birth before getting pregnant again. This is because shorter gaps between pregnancies raise the risk of [premature birth](#) and smaller babies. This increase applies to women of all ages, but the risk is highest among women aged 20-34.

For women over the age of 35, becoming pregnant less than 12 months after giving birth also increases the chance of serious complications for the mother. For women under 35, having babies close together does not increase the risks to their health.

Folic acid supplements

You should take [folic acid tablets \(supplements\)](#) if you are thinking about becoming pregnant. Ideally, do this from at least one month before you get pregnant; continue taking them until at least the end of the 12th week of pregnancy. This is even if you are healthy and have a good diet.

Folic acid is a vitamin which occurs naturally in certain foods such as spinach, sprouts, broccoli, green beans and potatoes. Some bread and breakfast cereals are fortified with folic acid. However, the intake for each individual can vary. When you are pregnant, you need a good supply of folic acid to help with the development of the baby. If you take folic acid tablets in early pregnancy you reduce the risk of having a baby born with a spinal cord problem such as [spina bifida](#).

Folic acid is a naturally occurring vitamin which your body needs. By taking these supplements you are making sure that you have a good, regular amount of folic acid which you need especially during pregnancy. You can buy folic acid tablets from pharmacies. Also, in the UK, [the NHS Healthy Start scheme](#) provides vitamin supplements that contain folic acid. These are free to many women who are on certain benefits.

For most women, the dose is 400 micrograms (0.4 mg) a day. If you have a higher risk of having a child with a spinal cord problem then the dose is 5 mg a day. You need a prescription for this higher dose - that is, if:

- You have had a previously affected pregnancy.
- You or your partner have (or have a family with) a spinal cord defect.
- You are taking [medication for epilepsy](#).
- You are obese - especially if your body mass index (BMI) is 30 or more.
- You have [coeliac disease](#) or [diabetes](#).
- You have [sickle cell anaemia](#), or [thalassaemia](#).

Ideally, start taking folic acid tablets before becoming pregnant. The common advice is to start from the time you start to plan to become pregnant. If the pregnancy is unplanned then start taking folic acid tablets as soon as you know that you are pregnant.

Because of the substantial benefits of folic acid, some countries routinely fortify staple foods - such as wheat, cornflour or rice - with folic acid. Currently there is debate as to whether the UK should follow suit and fortify certain common foods with folic acid.

Vitamin D supplements

Vitamin D is needed for growth. Vitamin D supplements are recommended for all pregnant women, breastfeeding women and breastfed babies. If you are at risk of having a low vitamin D level, it may be worth starting to build up the level in your body *before* you become pregnant. The following might make you more at risk of vitamin D deficiency.

- We obtain much of our vitamin D from sunlight, so you might be more at risk of low levels if you do not have much exposure to sunlight. For example, if you only go out covered up or if you are housebound.
- If you are of South Asian, African, Caribbean, or Middle Eastern family origin.
- If your diet is low in vitamin D. For example, if you eat no oily fish, eggs, meat, or vitamin D-fortified margarine or breakfast cereal.
- If you are overweight, with a [BMI](#) which is more than 30.

You can buy supplements from a pharmacy. [See also the separate leaflet called Vitamin D Deficiency for more information.](#)

Alcohol

Before becoming pregnant, consider how much alcohol you drink. Once pregnant, the advice will be to avoid all alcohol, at least for the first three months, and ideally throughout your pregnancy. So if you are planning a pregnancy, it is wise to cut down your alcohol intake and ideally to stop altogether. If you do this, you won't find you have been drinking alcohol before you realised you were pregnant. [See the separate leaflet called Diet and Lifestyle during Pregnancy for more information](#) about the possible damage alcohol can cause a developing baby.

If you think you are dependent on alcohol and may not be able to stop drinking without help then see your doctor for advice. [See also the separate leaflet called Alcohol and Sensible Drinking for more details.](#)

Smoking

Smoking can be harmful to your baby, as well as to you, so should you become pregnant, you are strongly advised not to smoke. See the separate leaflets called [Pregnancy and Smoking](#) and [Quit Smoking \(Smoking Cessation\)](#) for more information about the harm smoking can cause. It may take some time to quit, however. So if you smoke, ideally it is worth working on this *before* you become pregnant. Also more options are open to you to help you give up smoking when you are not pregnant. [See the separate leaflet called How to Quit Smoking for more information.](#)

Recreational drugs

[If you take or inject recreational \(illicit\) drugs](#), you are strongly advised to stop before becoming pregnant. It is best to postpone the pregnancy until you are off drugs. If you have an unplanned pregnancy you should aim to stop the drugs as soon as possible. If you are unable to do so, see your doctor for advice and a referral to a community drug team. [See the separate leaflet called Diet and Lifestyle during Pregnancy to read more about the effects recreational drugs can have in pregnancy](#), and how the community drug teams can help.

Overweight and obesity

If you are [overweight or obese](#), try to [lose some weight](#) before becoming pregnant. Women who are overweight or obese have more difficulty conceiving. There are also more likely to be complications in pregnancy if you are overweight, such as:

- [Diabetes in pregnancy.](#)
- [High blood pressure in pregnancy.](#)
- Blood clots (thromboembolism) in pregnancy.
- Needing a [caesarean section](#).
- [Premature](#) deliveries.
- Babies with abnormalities such as [spina bifida](#).

The more overweight you are, the greater the risk. Being underweight is also associated with problems in becoming pregnant and complications in pregnancy. If your [BMI is between 18.5 and 24.9](#), you have ideal body weight in preparation for pregnancy.

Food and diet

Generally speaking, you should eat a normal healthy balanced diet if you are planning to become pregnant. There is some special advice for pregnant women about foods and drinks to avoid. You may wish to refer to this so that if you think you might be pregnant, you can start to follow this advice straightaway. [See the separate leaflet called Diet and Lifestyle during Pregnancy.](#)

Medication

The effects of some prescribed medicines have been well studied and it is known that certain medicines are safe in pregnancy. For example, paracetamol at normal dose is

safe and useful for headaches, backache and other aches and pains that may occur during pregnancy.

However, some medicines are not safe and may be harmful to a developing baby - in particular, if you take them in the early weeks of pregnancy. Therefore, always tell a doctor or dentist who prescribes medication for you that you intend to become pregnant. Also, don't take medicines that you can buy over the counter (including herbal remedies) unless they are known to be safe in pregnancy. The pharmacist will advise you about this. Always tell a doctor or dentist who prescribes your medication that you intend to become pregnant.

If you already take regular medication, it is extremely important that you discuss this with a doctor before becoming pregnant. It may be that this medication is not safe for use in pregnancy and you will be changed to a safer alternative. In some cases you will be referred to a specialist to discuss this. Medication which may need to be changed includes:

- [Medication for epilepsy.](#)
- [Medication for diabetes type 1 or type 2.](#)
- Medication for mental health conditions such as [depression](#), [bipolar disorder](#) or [schizophrenia](#).
- Medication for [high blood pressure](#), particularly pills which are [angiotensin-converting enzyme \(ACE\) inhibitors](#) or angiotensin receptor blockers such as [losartan](#) or [candesartan](#).
- Medication for high cholesterol, such as [statins](#).

If you are on any medication which needs changing, continue using [contraception](#) until this has happened. This is so that you do not become pregnant whilst on medication which may harm your baby. If you have an unplanned pregnancy, discuss with your doctor as early as possible any medication that you take. In some cases, the risk of taking the medicine has to be balanced against the risk of not taking the medicine and your condition not being treated.

Infections

Rubella (German measles)

If you plan to become pregnant for the first time, you should check that you are immune to [rubella before becoming pregnant](#). See your practice nurse for a pre-pregnancy blood test. Most women are immune to rubella, as they have been immunised as a child. However, childhood immunisation does not work in every child and you may not be immune. [If you are not immune, you can be immunised.](#)

Note: you should not become pregnant for one month after the injection and ideally until your immunity has been confirmed by a further blood test.

The rubella virus causes a mild illness but can seriously damage an unborn baby, especially in the early stages of growth. So, until you know that you are immune (from the result of the blood test), you should avoid anyone who has rubella, especially in the first 16 weeks of pregnancy.

Chickenpox

[Having chickenpox when pregnant](#) can be a nasty illness and there is some risk to the developing baby. A vaccine is offered to healthcare workers (doctors, nurses, etc) who have not previously had chickenpox and so are not immune and may catch chickenpox. (About 1 adult in 10 has not had chickenpox as a child.) Non-immune healthcare workers should consider having this vaccination before becoming pregnant. If you are not sure if you have had chickenpox, a blood test can check if you have previously had it.

Hepatitis B

A mother who is [infected with hepatitis B](#) has a high risk of passing it on to her newborn baby. If you are at high risk of catching hepatitis B, you should be [immunised against this virus](#) before becoming pregnant. People at increased risk and who should be immunised include:

- Those whose job puts them at risk of contracting hepatitis B - for example, healthcare personnel and staff at daycare or residential centres.
- Those who inject recreational (illicit) drugs.
- Those who change sexual partners frequently.
- Those who live in close contact with someone infected with hepatitis B.

Consider your working environment

If you think that your job may pose a risk to a pregnancy then ideally you should discuss this with your employer before becoming pregnant. Certain jobs and workplaces may pose a risk to a pregnancy, in particular to the early stages of pregnancy. For example:

- If you work with raw meat. Raw meat is sometimes contaminated with germs, such as listeria and toxoplasma. If these germs infect adults, they may cause [listeriosis](#) or [toxoplasmosis](#), but may cause little harm. However, these germs can cause serious problems to your unborn child if you become infected when you are pregnant.
- If you work with certain animals. For example:
 - You should avoid contact with sheep and lambs at lambing time. This is because some lambs are born contaminated with germs such as listeria, toxoplasma and chlamydia, which may affect you and your unborn baby.
 - Cats and kittens often carry toxoplasma germs - especially in cat poo (faeces). So, cleaning out cat litter trays and handling cats and kittens can be a risk.
- If your job puts you at risk of contracting [hepatitis B](#), you should be immunised against this virus. For example, if you are a healthcare worker, or work at a daycare or residential centre. (Women with hepatitis B can pass on the infection to their child at childbirth.)
- If you work with chemicals, fumes, radiation, etc. Some may be toxic to an unborn baby.

The above are just examples. In short, if you think that your occupation may pose a risk to a pregnancy then you should discuss this with your employer before becoming pregnant. A change in job, or in working practice, may be necessary.

You and your employer may wish to download [information guides from the Health and](#)

[Safety Executive \(HSE\) website](#). They have an information line if you have a health and safety concern at work but do not wish to discuss with your employer your intention to become pregnant.

Medical conditions

Women with certain medical conditions may benefit from advice **before** becoming pregnant. If you have any of the conditions listed below, it would be sensible to continue using [contraception](#) until you have discussed with your doctor that you are considering becoming pregnant. If you have an unplanned pregnancy and have any of these conditions, see your doctor as soon as possible. Do not stop any medication without talking with your doctor.

Diabetes

If you have diabetes, it is extremely important to see your doctor BEFORE you become pregnant. This is because good control of your sugar levels, both before pregnancy and in early pregnancy, can reduce various risks. Diabetes increases your risk of [miscarriage](#) and stillbirth. It also increases the risk that your baby will be born with abnormalities, or become seriously ill after birth. However, good control of sugar levels starting before pregnancy can reduce these risks. You will be referred to a specialist in diabetes, and your medication (and/or insulin) may be changed. You will have checks on your eyes and your kidneys. You will work with your specialist together to get your blood glucose levels within a very tight range. The better the control before pregnancy and during early pregnancy, the better the outcome. This is why you should plan ahead and see your GP or specialist before you get pregnant. [See the separate leaflet called Diabetes and Pregnancy.](#)

Epilepsy

Many treatments for epilepsy are not safe to take in pregnancy. See your doctor or epilepsy specialist to discuss this BEFORE becoming pregnant. You may need to be switched to another medication. However, it is risky to have fits in pregnancy too, so do not stop your medication suddenly without discussing it with your specialist. It is advisable to keep taking [contraception](#) until you have had this discussion. [See the separate leaflet called Epilepsy and Planning Pregnancy](#)

High blood pressure

If you are on [medication for high blood pressure \(hypertension\)](#), ideally see your GP before becoming pregnant. As discussed in the medication section above, there are some blood pressure tablets which should not be taken if you are pregnant. Your doctor will advise if you need to change and may refer you to a specialist for further advice.

Mental health

If you are taking medication for [depression](#), [anxiety](#), [bipolar disorder](#) or [schizophrenia](#), see your doctor to discuss this before becoming pregnant. You and your GP or mental health specialist will need to weigh up the risks of taking medication in pregnancy against the risks of becoming mentally ill if you stop. It may be that your medication will

be changed to be safer in pregnancy. Unfortunately, there is not much information available on how safe most medication is in pregnancy. Your doctor will explain what is known about your medication, and together you can make a decision about whether you need to stop or change.

If you have a past history of severe mental health illness, it is best to be under the care of a specialist who can monitor you during pregnancy. You should have access to details about who you should contact if you feel you are becoming unwell.

Hereditary conditions

If your family has a history of a condition that runs in families, you may wish to have [genetic testing](#) before planning a pregnancy. This will help work out your chances of having that condition. Examples of conditions which may run in families include:

- [Huntington's disease](#).
- Neurofibromatosis.
- Tuberous sclerosis.
- [Marfan syndrome](#).
- Adult polycystic disease.
- Achondroplasia.
- [Cystic fibrosis](#).
- Tay-Sachs disease.
- Gaucher's disease.
- Friedreich's ataxia.
- [Congenital adrenal hyperplasia](#).
- Spinal muscular atrophy.
- [Duchenne muscular dystrophy](#).
- Fragile X syndrome.
- Haemophilias A and B.
- Glucose-6-phosphate dehydrogenase deficiency.

Other medical conditions

Other conditions for which it is best to see a doctor before becoming pregnant include:

- Thyroid problems - [an underactive thyroid gland \(hypothyroidism\)](#) or [an overactive thyroid gland \(hyperthyroidism\)](#).
- Kidney disease, such as [chronic kidney disease](#).
- [Rheumatoid arthritis](#), especially if you are on medication.
- If you have ever had [deep vein thrombosis](#) or [pulmonary embolism](#), whether or not you are currently on treatment.
- Heart disease.
- Blood conditions such as [sickle cell anaemia](#) or [thalassaemia](#).

Screening tests

Pregnant women in England, Scotland and Wales and those at high risk in Northern Ireland are offered a screening test for [sickle cell disease](#) and [thalassaemia](#). Some women may wish to have the screening test before becoming pregnant, especially if their family origins make these diseases more likely. If you or your partner or either of your families originate from Africa, the Caribbean, Southeast Asia, Sardinia, Greece, Turkey or Cyprus, it may be worth having a screening test before pregnancy. See your

doctor to discuss this. Genetic testing is a rapidly developing area of medicine. It may be possible that more tests will become available to detect carriers of various diseases. When they become available, these may be tests to consider before becoming pregnant.

Cervical smear

All women from the age of 25 should have a [three-yearly cervical smear](#) to check for early pre-cancer changes to the neck of the womb (cervix). (In the UK women are invited every three years between the ages of 25 and 49, and every five years between the ages of 50 and 64.) This cannot be done in pregnancy or for three months after a baby is born. So, if you are thinking of becoming pregnant, check with your doctor or nurse to see if your smear is due. If it is due, have it before becoming pregnant.

Summary and checklist

Most pregnancies go well and without any major problems. But, it is wise to reduce any risks as much as possible. So, below is a reminder of things to consider before becoming pregnant, and as soon as you realise you are pregnant:

Things you should do now

- See your doctor to discuss pregnancy if you have any long-standing medical condition or take any regular medication. Even if you do not, it may be worth visiting your doctor or practice nurse to check your cervical smear is up to date and to have a blood test to check to see if you are immune to [rubella](#) (German measles). Your doctor or nurse will advise if any other tests are needed.
- If you have [diabetes](#) or are on medication for [epilepsy](#) see a doctor sooner rather than later; continue contraception until any plans to control your medical condition are in place.
- Take folic acid tablets before you become pregnant and until 12 weeks of pregnancy.
- Eat a healthy diet.

Things you should stop or cut down on

- Alcohol - you are strongly advised not to drink at all.
- Smoking - you are strongly advised to stop completely.
- Street (illicit) drugs - you are strongly advised to stop completely.

Other things to consider

- Immunisation against hepatitis B if you are at increased risk of getting this infection.
- Immunisation against chickenpox if you are a healthcare worker and have not previously had chickenpox and so are not immune.
- Your medication - including herbal and 'over-the-counter' medicines. Are they safe?
- Your work environment - is it safe?
- Medical conditions in yourself, or conditions which run in your family.

- Screening tests for [sickle cell disease](#) and [thalassaemia](#).

Further reading and references

- [Antenatal care for uncomplicated pregnancies](#); NICE Clinical Guideline (March 2008 - updated February 2019)
- [Feldman HS, Jones KL, Lindsay S, et al](#); Prenatal alcohol exposure patterns and alcohol-related birth defects and growth deficiencies: a prospective study. Alcohol Clin Exp Res. 2012 Apr;36(4):670-6. doi:
- [Diabetes in pregnancy - management from preconception to the postnatal period](#); NICE Clinical Guideline (February 2015 - last updated December 2020)
- [De-Regil LM, Pena-Rosas JP, Fernandez-Gaxiola AC, et al](#); Effects and safety of periconceptional oral folate supplementation for preventing birth defects. Cochrane Database Syst Rev. 2015 Dec 14;12:CD007950. doi: 10.1002/14651858.CD007950.pub3.
- [UK Chief Medical Officers' Low Risk Drinking Guidelines](#); GOV.UK, August 2016
- [Valproate use by women and girls](#); Department of Health, GOV.UK

Article History

The information on this page is written and peer reviewed by qualified clinicians.

- Next review due: 27 Apr 2025
- 28 Apr 2020 | Latest version
Last updated by
[Dr Hayley Willacy, FRCGP](#)
- Peer reviewed by
[Dr Colin Tidy, MRCGP](#)

ARTICLE INFORMATION

LAST REVIEWED	28 April 2020
Next Review	27 April 2025
Document ID	4240 (v49)
Author	Dr Hayley Willacy
Peer reviewer	Dr Colin Tidy

Disclaimer: This article is for information only and should not be used for the diagnosis or treatment of medical conditions. Egton Medical Information Systems Limited has used all reasonable care in compiling the information but make no warranty as to its accuracy. Consult a doctor or other health care professional for diagnosis and treatment of medical conditions. For details see out **conditions**